

Inverse problems of heat conduction...

27554
S/170/61/004/010/008/019
B109/B138

4

There are 2 Soviet references.

ASSOCIATION: Tekhnologicheskii institut rybnoy promyshlennosti i
khozyaystva, g. Kaliningrad (Technological Institute of the
Fishing Industry and Economy, Kaliningrad)

SUBMITTED: March 25, 1961

Card 9/9

VINER, A.M., inzh.; TEMKIN, A.G., kand.tekhn.nauk; FEDOROV, V.N., inzh.

Nomogram for calculating heat transfer in a furnace. Teploenergetika
8 no.1:89-90 Ja '61. (MIRA 14:4)
(Furnaces) (Heat--Transmission)

S/170/62/005/004/012/016
B104/B108

AUTHOR: Temkin, A. G.

TITLE: The temperature field of bodies of complex shape at regular heat conditions

PERIODICAL: Inzhenerno-fizicheskiy zhurnal, v. 5, no. 4, 1962, 106 - 121

TEXT: General properties of body shape criteria (characteristic numbers), the temperature field near the center of a body, the mean distance between isotherms, the heat conduction equation, and errors found in various publications are discussed in the present survey which is based on publications, issued between 1933 and 1961. N. Ye. Zhukovskiy, A. V. Lykov, A. S. Ginzburg, D. V. Burdin, V. A. Krasovskiy, G. Ye. Pikus, L. S. Leybenzon, G. P. Buynyachenko, I. A. Parkhomenko, L. V. Kravchuk, G. M. Kondrat'yev are mentioned. There are 43 references: 42 Soviet and 1 non-Soviet.

ASSOCIATION: Tekhnicheskiy institut rybnoy promyshlennosti i khozyaystva, g. Kaliningrad (Technical Institute of Fish Industry and Fisheries, Kaliningrad)

~~Card 1/2~~

TEMKIN, A.G.

Apropos of A.N. Borshchevskii's article "Heat transfer
through enclosing structures." Inzh.-fiz. zhur. 5 no.6:118-120
Je '62. (MIRA 15:12)
(Heat—Transmission) (Structures, Theory of)
(Borshchevskii, A.N.)

41316

S/170/62/005/010/009/009

B104/B186

26.5/100

AUTHOR: Temkin, A. G.

TITLE: Temperature field of a multilayered wall

PERIODICAL: Inzhenerno-fizicheskiy zhurnal, v. 5, no. 10, 1962, 104 - 117

TEXT: If k finite measurement values are available the equation of heat conduction through a multilayered wall is

$$C(r) \frac{\partial t}{\partial \tau} = r^{1-k} \frac{\partial}{\partial r} \left[\lambda(r) r^{k-1} \frac{\partial t}{\partial r} \right] \quad (1.1).$$

The specific heat C and the coefficient of heat conduction λ of this wall depend considerably on the coordinates and are piece-wise continuous functions. ✓

$$\frac{\partial t}{\partial F} = \frac{1}{C(N) N^{k-1}} \frac{\partial}{\partial N} \left[\Lambda(N) N^{k-1} \frac{\partial t}{\partial N} \right] \quad (1.6)$$

is obtained from (1) by introducing the dimensionless coordinate $N = r/r_e$, the Fourier number $F = \alpha_0 \tau / r_e^2$, the dimensionless thermal conductivity

Card 1/4

Temperature field of a multilayered...

S/170/62/005/010/009/009
B104/B186

$\Lambda(N) = \lambda(r)/\lambda_0$ and the dimensionless specific heat $C(N) = C(r)a_0\lambda_0$.
 $t(N_1, F) = t_1(F)$, $\Lambda(N_2) \frac{\partial t(N, E)}{\partial N} = q(F)$ are the mixed boundary conditions.

The solution is obtained as a series

$$t(N, F) = t_1(F)1 + t_2'(F)\nabla^{-1}1 + \dots + t_1^{(n)}(F)\nabla^{-n}1 + \dots + \\ + q(F)a(N) + q'(F)\nabla^{-1}a + \dots + q^{(n)}(F)\nabla^{-n}a + \dots \quad (2.4)$$

arranged with respect to the derivatives of the quantities to be measured and the radial quasipolynomials of the problem. The function (2.4) describes the temperature field at sufficiently long time values for the initial temperature distribution no longer to influence the heat conduction. The Fourier integral makes it possible to construct the field of the aftereffect of this problem. It can be shown that this field gradually vanishes and that at the initial moment a temperature distribution appears contradicting the initial value of the function (2.4). If the temperatures on the surfaces N_1 and N_2 are known functions of time,

then the field of the action can be represented as the sum of the two series.

Card 2/4

Temperature field of a multilayered...

S/170/62/005/010/009/009
B104/B186

$$t(N, F) = \sum_{n=0}^{\infty} t^{(n)}(F) P_n(N, N_1) + t^{(n)}(F) P_n(N, N_2), \quad (3.3).$$

The series are arranged accordingly with respect to the derivatives of the temperatures to be measured and with respect to the radial quasipolynomials of the problem. The convective heat exchange with media at variable temperatures is studied with the aid of the boundary conditions

$$[t_1(F) - t(N_1, F)] B_1 = -\Lambda(N_1) \frac{\partial t(N_1, F)}{\partial N} \quad (4.1)$$

$$[t(N_2, F) - t_2(F)] B_2 = -\Lambda(N_2) \frac{\partial t(N_2, F)}{\partial N} \quad (4.2).$$

of the third kind. For the field of action

$$t(N, F) = t_1(F) P_0(N, N_1) + t'_1(F) P_1(N, N_1) + \dots + t^{(n)}(F) P_n(N, N_1) + \dots \\ + t_2(F) P_0(N, N_2) + t'_2(F) P_1(N, N_2) + \dots + t^{(n)}(F) P_n(N, N_2) + \dots, \quad (4.4)$$

is obtained in a similar way. This series is arranged with respect to the
Card 3/4

Temperature field of a multilayered...

S/170/62/005/010/009/009
B104/B186

derivatives of the temperatures of the inner and outer medium. If the temperature on the inner surface is kept constant, and if the temperature on the outer surface varies periodically, then

$$t(N, F) = t_1 P_0(N, N_1) + \\ + t_0 [P_0(N, N_2) - \omega^2 P_2(N, N_2) + \omega^4 P_4(N, N_2) - \dots] \cos \omega F + \\ + t_0 [-\omega P_1(N, N_2) + \omega^3 P_3(N, N_2) - \omega^5 P_5(N, N_2) + \dots] \sin \omega F, \quad (5.20)$$

is valid for the field of action.

ASSOCIATION: Tekhnicheskii institut rybnoy promyshlennosti i khozyaystva,
g. Kaliningrad (Technical Institute of the Fish Industry and
Fisheries, Kaliningrad)

SUBMITTED: July 23, 1962

Card 4/4

TEMKIN, A.G.

Phenomenological equation of transfer processes. Dokl. AN BSSR
7 no.2:92-94 F '63. (MIRA 16:7)

1. Tekhnicheskii institut rybnoy promyshlennosti i khozyaystva,
Kaliningrad. Predstavleno akademikom AN BSSR A.V. Lykovym.
(Heat-Transmission) (Mass transfer)

TEMKIN, A.G.

Solution of a Schrodinger type equation. Dokl. AN BSSR 7
no.4:240-243 Ap '63. (MIRA 16:11)

1. Energeticheskiy institut AN BSSR. Predstavleno akademikom
AN BSSR A.V. Lykovym.

TFMKIN, A.G., kand.tekhn.nauk, dotsent; IONOV, A.G., aspirant

Useful book on refrigerating engineering. Khol.tekh. 40 no.6:
54-55 N-D '63. (MIRA 17:4)

$\frac{1}{2} \pi (H_1 + H_2) \cdot \frac{1}{2} \pi (H_1 + H_2)$

Reconstruction of the fields of heat and mass contents of a colloidal capillary-porous body. *Inzh.-fiz. zhurn.* no. 10:86-93 0 164. (MIRA 10:11)

1. Tekhnicheskiiy institut rybnoy promyshlennosti i khozyaystva.
Kuliningrad.

L 3639-66 EMT(1)/EPF(c)/ETC/EPF(n)-2/ENG(n) WW
ACCESSION NR: AP5022384

UR/0170/65/009/003/0305/0317
536.75+536.24

55
52
B

AUTHOR: Temkin, A. G. 44, 55

TITLE: Determination of the parameters of internal heat and mass transfer using the characteristic functions of the thermodynamics of irreversible processes

SOURCE: Inzhenerno-fizicheskiy zhurnal, v. 9, no. 3, 1965, 305-317

TOPIC TAGS: heat transfer 21, 44, 55, mass transfer, thermodynamics, irreversible process, mathematic matrix, thermal diffusion

ABSTRACT: The article presents a theoretical development starting with the determination of the parameters of internal transfer first with multiple point changes in temperature and then with two point changes in temperature. The methods of matrix mathematics are applied to a determination of transfer criteria using a local measurement of the moisture content. It is demonstrated that, starting from any given set of experimentally determined parameters, the rest can be determined by use of the characteristic thermodynamic functions of irreversible processes. This includes the coefficients of thermal diffusivity and

Card 1/2

L 3639-66

ACCESSION NR: AP5022384

the criteria for inertial and thermal diffusion and phase transfer. Examples of
typical calculations by this method are presented in tabular form. Orig. art. has
74 formulas and 3 tables

ASSOCIATION: Politeknicheskiy institut, g. Riga (Polytechnical Institute, Riga)

SUBMITTED: 00

ENCL: 00

SUB CODE: TD

NR REF SOV: 019

OTHER: 002

44,55

BVR.
Card 2/2

Russia (1923) Manual on planning prestressed reinforced concrete structures (A-148-52)
TA444.RB MSPTI

1. Concrete, Prestressed. I. Temkin, A. I., ed. II. Moscow. Tsentral'naya nauchno-issledovatel'skii institut promyslennykh sooruzhenii.

LENNIN, A. I., 22.

Manual on planning prestressed reinforced concrete structures (I-148-52)
MSPTI

Razrabotana TSentr. nauchno-issledovatel'skim in-tom promyshl. sooruzhenii. Utverzhdena
10 okt. 1952 g. Moskva, Gos. izd-vo lit-ry po stroitel'stvu i arkhitekture, 1953.
81 p. (54-33025)

TA444.R8

I. Concrete, Prestressed. II. Moscow. TSentral'nyi nauchno-issledovatel'skii institut
promyshlennykh sooruzhenii.

NEVYAZHSKIY, I.Kh; DRABKIN, V.F.; TRUBETSKOY, V.F.; TEMKIN, A.S.

Use of ferrite-core inductance in the high-frequency power stage
circuit of the proton synchrotron. Radiotekh.i elektron.i no.7:954-
964 J. '56. (MIRA 10:1)

(Synchrotron)

h6748

S/120/62/000/004/016/047
E192/E382

24.6730

AUTHORS: Lebedev-Krasin, Yu.M., Gutner, B.M., Pisarevskiy, V.Ye.,
Temkin, A.S., Barabash, L.Z., Kuryshev, V.S. and
Moiseyev, A.I.

TITLE: The accelerating elements of the proton synchrotron
and the system of their high-frequency feed

PERIODICAL: Priory i tekhnika eksperimenta, no. 4, 1962,
94 - 97

TEXT: The description, principal characteristics and the
results of the control of the h.f. accelerating system of the
7 GeV proton cyclotron are reported. The accelerating elements
are in the form of drift tubes situated in 11 compensating
magnets. Each of the 11 electrodes is fed from a separate
system of high-frequency amplifiers consisting of a 7-stage
wideband amplifier and an automatically-tuned resonance output
amplifier. The inductances of the resonant circuit in the output
stages are in the form of coils fitted with ferrite cores. The
amplitude of the high-frequency field of each accelerating
electrode is 2.5 kV \pm 10% over the frequency range of
Card 1/2

The accelerating elements

S/120/62/000/004/016/047
E192/E382

0.65 - 8.5 Mc/s. The phase-shift between the output voltages of any two channels is less than 30° . The overall power used by the supply system is 400 kVA. By using tuned amplifiers in the output stages the power consumption was reduced by about 30 times, as compared with a non-tuned amplifier. There are 4 figures.

SUBMITTED: March 29, 1962

Card 2/2

I 3778-66 EWT(m)/EWA(m)-2 IJP(c) GS
ACCESSION NR: AT5007965

S/0000/64/000/000/0932/0936

AUTHOR: Vodop'yanov, F. A.; Zhukovskiy, L. S.; Zaimanov, V. B.; Ivanov, Yu. S.;
Izergina, Ye. V.; Kuz'min, A. A.; Prokop'yev, A. I.; Temkin, A. S.; Rubchinskiy,
S. M.

TITLE: System for the generation of the accelerating field of a 70-Gev proton
synchrotron /9.

SOURCE: International Conference on High Energy Accelerators. Dubna, 1963.
Trudy. Moscow, Atomizdat, 1964, 932-936

TOPIC TAGS: high energy accelerator, synchrotron, particle beam, magnetic field

ABSTRACT: After the development of a high-precision system of frequency control of the accelerating field of the proton 50-60 Gev synchrotron with critical energy compensation (Mints, A. L., et al., Proc. International Conference on High Energy Accelerators and Instruments, CERN 1959), it was decided to achieve an alternative accelerator with transition through the critical energy, which makes it possible to increase the energy to 70 Gev. In this modification of the accelerator serious difficulties are encountered with the realization of a system for generating an accelerating field with frequency control only according to the N-program. Therefore,

Card 1/3

1. 3778 66

ACCESSION NR: AT5007965

it was decided to achieve a system with twin frequency control: rough, according to the H -program, and precise, according to the information on the radial and phase position of the accelerated particle beam. The present report discusses the principal characteristics governing the achievement of a programmed FM-generator, a system of frequency control according to information of the position of the accelerated particle bunches, and accelerator installation. The programmed FM-generator consists of the usual elements: transducer of the derived magnetic field strength (inductive coil in the gap of the measuring electromagnet), electronic switch, tube integrator, modulator, FM-oscillator, phase manipulator, amplitude modulator of accelerating voltage, amplifier-distributor, and a system of cable contacts. To obtain energy increase per revolution of $\Delta E = 166$ Kev for a rate of change of magnetic field strength of $\dot{H} = 550$ oersteds/second and $\phi_s = 30^\circ$, provision is made for the application of 53 accelerator stations with rated input of 7 kilovolts and 6 kilowatts power. Provisions are also made for the short-duration increase of this voltage, 1.8 times up to the time of beam bunching (around 15 microseconds), and its slow decrease to about 2 times less toward the end of the acceleration cycle with the aim of preserving constant equilibrium phase during the fall in the magnetic field growth rate. The system of frequency control of the accelerating field according to the information on the accelerated particle beam position is similar in

Card 2/3

L 3778-66

ACCESSION NR: AT5007965

principle of operation to a system described by Yu. S. Ivanov and A. A. Kus'min (Priory i tekhnika eksperimenta, No. 4, 106, (1962)), which was intended to stabilize the position of the center of gravity of the beam according to radius and phase. Orig. art. has: 1 figure.

ASSOCIATION: Radiotekhnicheskiy institut AN SSSR (Radio Engineering Institute, AN SSSR)

SUBMITTED: 26May64

ENCL: 00

SUB CODE: NP 12

NO REF SOV: 001

OTHER: 001


Card 3/3

NESTEROV, S.N.; VALETOV, V.V., inzhener, redaktor; TEMKIN, A.V., redaktor;
GENICH, V.A., kandidat tekhnicheskikh nauk, retsenzent; UVAROVA,
A.F., tekhnicheskii redaktor.

[Establishing norms for use of materials in machine building
plants; method of determining consumption rates of basic and
subsidiary materials for plants engaged in mass and large-scale
production] Normirovanie raskhoda materialov na mashinostroi-
tel'nykh zavodakh; metodika opredeleniia norm raskhoda osnovnykh
i vspomogatel'nykh materialov na zavodakh massovogo i krupno-
seriinogo proizvodstva. Moskva, Gos.nauchno-tekhn.isd-vo
mashinostroit. lit-ry, 1955. 187 p. [Microfilm] (MLRA 8:12)
(Machinery industry)

TEMKIN, A.V.

BARANOV, A.I., kandidat tekhnicheskikh nauk; KUZ'MIN, V.V., inzhener;
GOKUN, V.B., kandidat tekhnicheskikh nauk, retsenzent; NOVIKOV, .
K.D., inzhener, retsenzent; TKACHENKO, V.V., kandidat tekhnicheskikh
nauk, redaktor; TEMKIN, A.V., redaktor; UVAKOVA, A.F., tekhnicheskii
redaktor; SOKOLOVA, I.F., tekhnicheskii redaktor.

[Setting standards and norms in machine building] Standartizatsiia i
normalizatsiia v mashinostroenii. Izd.2-oe, perer. i dop. Moskva,
Gos.nauchno-tekhn.izd-vo mashinostroitel'noi lit-ry, 1955. 202 p.
(Machinery industry) (MLRA 8:11)

SHEVELEV, M.L.; TIKHONOV, A.S., kandidat tekhnicheskikh nauk dotsent, retsenzent;
SHAROV, N.V., inzhener, retsenzent; PCHELINTSEV, V.A., inzhener, retsenzent;
TEMKIN, A.V., redaktor; MATVEYEVA, Ye.N., tekhnicheskiiy redaktor.

[Fire prevention in machine building] Protivopozharnaya tekhnika v
mashinostroyeni. Izd. 2-Oe, perer. i dop. Moskva, Gos. nauchno-tekhn.
izd-vo mashinostroyit. lit-ry, 1955. 208 p. (MIRA 9:6)
(Factories--Fires and fire prevention)

ASVAL'DOV, M. Ya.; TEMKIN, A. V., redaktor; UVAROVA, A. F., tekhnicheskii
redaktor

[Complex improvement in the operation of metal cutting machines]
Kompleksnoe uluchshenie ispol'zovaniia metalloreshushchikh stankov.
Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry, 1955.
229 p. (MLRA 9:2)

(Cutting tools)

ZUBOK, V.N., inzhener, redaktor; UMYAGIN, M.G., inzhener, redaktor;
KASSATSIER, M.S. inzhener, redaktor; SHIFRIN, S.M., redaktor;
TEMKIN, A.V., redaktor; TIKHONOV, A. Ya., tekhnicheskii redaktor.

[Experience in introducing advanced technology in factories
engaged in heavy machine building] Opyt vnedreniia peredovoi tekhnologii na zavodakh tiashelogo mashinostroeniia. Moskva, Gosnauchno-tekhn.izd-vo mashinostroit.lit-ry, 1955.306 p. (MLRA 9:4)

1.Moscow. Vsesoiuznyy proyektno-tekhnologicheskii institut.
(Machinery--Construction)

IVANOV, Nikolay Filippovich; GORINSHTEYN, B.I., retsehsent; BYKHENVAL'D, A.V.,
kandidat ekonomicheskikh nauk, dotsent, redaktor: ~~TRUKHIN, A.V.~~
redaktor izdatel'stva; POPOVA, S.M., tekhnicheskii redaktor

[Operational planning; planning machine inspection every ten days
at machine building plants producing in lots] Operativnoe planiro-
vanie; podskadnoe, mashinokomplektnoe planirovanie na mashino-
stroitel'nykh zavodakh seriinogo proizvodstva. Moskva, Gos. nauchno-
tekh. izd-vo mashinostroit. lit-ry, 1956. 105 p. (MLRA 10:3)
(Machinery industry)

GANSHTAK, V.I.; BOGINSKIY, I.N., inzhener, redaktor; TEMKIN, A.V., redaktor;
UVAROVA, A.P., tekhnicheskii redaktor.

[Cost of production in machine-building] Sebestoinost' produktsii v
mashinostreemii. Izd. 2-ee, perer. i dop. Moskva, Gos.nauchno-tekhn.
izd-vo mashinostreitel'noi lit-ry, 1956. 153 p. (MIRA 9:6)
(Machinery industry--Costs)

Tom IV, 1956
IVANOV, Nikolay Vasil'yevich; MALYUTIN, Nikolay Kuz'mich; FLEYSHMAN, Abram
L'vovich; BURSHTAYN, I.I., retsenzent; LOBODIN, P.V., retsenzent;
MOROZOV, A.N., retsenzent; LYUBOVICH, Yu.O., kandidat ekonomicheskikh
nauk, redaktor; TEMKIN, A.V., redaktor izdatel'stva; UVAROVA, A.F.,
tekhnicheskiiy redaktor.

[Supply of materials and equipment in machinery manufacturing] Material'-
no-tekhnicheskoe snabzhenie v mashinostroenii. Moskva, Gos.nauchno-
tekhn.izd-vo mashinostroit.lit-ry, 1956. 275 p. (MIRA 10:4)
(Machinery industry)

BOCHAROV, Grigoriy Grigor'yevich; YUR'YEV, N.M., Inzhener, retsentyent;
SHNAYVAS, P.Kh., redaktor; TMMKIN, A.V., redaktor; EL'KIND, V.D.,
tekhnicheskiy redaktor

[Evaluating and calculating production in the machinery industry]
Uchet proizvodstva i kal'kulatsiya v mashinostroenii. Izd. 2-oe.
perer. Moskva, Gos.nauchno-tekhn.izd-vo mashinostroit. lit-ry,
1957. 309 p. (MLRA 10:10)
(Machinery industry)

ANDREYEV, Yevgeniy Dmitriyevich; GORENSHTEYN, B.I., retsenzent; KUZNETSOV, B.R., retsenzent; TEMKIN, A.V., red.; SALYANSKIY, A.A., red.izd-va; UVAROVA, A.F., tekhn.red.

[Operational and production planning in machinery plants with piece and small-scale production; organization by work schedules]
Operativno-proizvodstvennoe planirovanie na mashinostroitel'nom zavode edinichnogo i melkoseriynogo proizvodstva; rabota po grafiku. Izd. 2., dop. Moskva, Gos. nauchno-tekhn.izd-vo mashinostroit. lit-ry, 1958. 218 p. (MIRA 12:2)
(Machinery industry)

21.6000

also 2209

S/076/60/034/011/012/024
B004/B064AUTHOR: Tëmkin, A. Ya. (Moscow)

TITLE: Mechanism of the Radiolysis of Solid Oxalic Acid

PERIODICAL: Zhurnal fizicheskoy khimii, 1960, Vol. 34, No. 11,
pp. 2503-2505

TEXT: The purpose of the present paper isto prove that the radiochemical processes in solids take place along tracks. Using the experimental data of Refs. 1,2 the following relation is written: $\nu(t) = (wv/a)\ln(1+aQt)(2)$. $\nu(t)$ is the concentration of the radicals; w is the number of tracks per unit volume and unit time; v is the volume of the track; $Q = n(t_0)$ is the initial condition for the density n of the radicals; and a is the coefficient of recombination. This equation is in good agreement with the data of Ref. 1. Since a long lifetime of the free radicals must be assumed for solid substances, the overlapping of the tracks was studied, and a correction $\chi(t)$ was calculated for equation (2). It is found to increase with t , so that saturation is achieved after a long time of irradiation. Thus,

Card 1/2

86782

Mechanism of the Radiolysis of Solid
Oxalic Acid

S/076/60/034/011/012/024
B004/B064

the radiochemical processes in solid oxalic acid proceed along tracks according to a radical mechanism. This is assumed to hold also for other solid substances which are neither conductors nor semiconductors. L. S. Polak, N. Ya. Chernyak, Yu. N. Molin, and A. T. Koritskiy are thanked for a discussion. [Abstracter's note: The experimental data on which the calculation is based are not given.] There are 2 Soviet references.

ASSOCIATION: Akademiya nauk SSSR. Institut neftekhimicheskogo sinteza
(Academy of Sciences USSR, Institute of Petrochemical
Synthesis)

SUBMITTED: February 19, 1959

45/11

Card 2/2

FILIPPOV, Yevgeniy Mikhaylovich. Prinimali uchastiye: GUBERMAN, SH.A.; LEYPUNSKAYA, D.I., nauchnyy sotr., red.; BESPALOV, D.F., nauchnyy sotr., red.; SREBRODOL'SKIY, D.M., nauchnyy sotr., red.; SHIMELEVICH, Yu.S., nauchnyy sotr., red.; TEMKIN, A.Ya., red.; MEDER, V.M., red. izd-va; PRUSAKOVA, T.A., tekhn. red.; MAKUNII, Ye.V., tekhn. red.

[Applied nuclear geophysics; use of sources of nuclear radiation in geology and geophysics] Prikladnaya yadernaya geofizika; primeneniye istochnikov yadernogo izlucheniya v geologii i geofizike. Pod obshchei red. L.S.Polaka. Moskva, Izd-vo Akad. nauk SSSR, 1962. 579 p. (MIRA 15:12)

1. Chlen-korrespondent Akademii nauk SSSR (for Filippov). 2. Institut geologii i razrabotki goryuchikh iskopayemykh (for Ley-punskaya, Bepalov, Srebrodol'skiy, Shimelevich). 3. Institut neftekhimicheskogo sinteza Akademii nauk SSSR (for Temkin). (Nuclear geophysics)

TEMKIN, A. Ya.

USSR/Nuclear Physics - Meson formation

FD 429

Card 1/1

Pub. 147-15/16

Author

: Temkin, A. Ya.

Title

: Conversion of two photons into a π^0 -meson, and the production of π^0 -mesons in the Compton effect

Periodical

: Zhur. eksp. i teor. fiz. 26, 645-646, May 1954

Abstract

: A letter to the editors. Computes the total cross-section of meson production, and finds it to be very small (10^{-46} cm²) even at 21-22 Bev. Refers to various related works, appearing in 1950-1951 of: A. M. Baldin and V. V. Mikhaylov (ZhETF 20, 1057, 1950); B. Ioffe, A. Rudik and I. Shmushkevich (DAN SSSR 77,403,1951); V. B. Berestetskii and I. Ya. Pomeranchuk (DAN SSSR 77,803, 1951). Acknowledges the interest of Yu. M. Shirokov in the present work.

Institution

:

Submitted

: October 15, 1953

TEMKIN, A. YA.

SUBJECT USSR / PHYSICS

CARD 1 / 5

PA - 1774

AUTHOR TEMKIN, A. YA.

TITLE On the Theory of the Slowing Down of Neutrons.

PERIODICAL Zhurn. eksp. i teor. fis, 31, fasc. 5, 893-895 (1956)

Issued: 1 / 1957

The integral of collisions between neutrons and the nuclei of the slowing down material can be written down as follows:

$$\sum_{\alpha} \int_0^u du' \int_{-1}^1 d\mu' \int_0^{2\pi} d\beta' f_{\alpha 0}(u-u') \frac{\lambda(u')}{\lambda_{\alpha}(u')} \delta(\mu_0 - \gamma_{\alpha}) \Psi(\vec{r}, u', \mu', \beta') = \sum_{\alpha} \hat{K}_{\alpha} \hat{B}_{\alpha} \Psi$$

$$\hat{K}_{\alpha} \equiv \int_0^u du' f_{\alpha 0}(u-u') (\lambda(u')/\lambda_{\alpha}(u')) \int_{-1}^1 d\mu' K_{\alpha}(\mu, \mu', u-u')$$

$$K_{\alpha} \equiv (1 - \mu^2 - \mu'^2 - \gamma_{\alpha}^2 - 2\gamma_{\alpha} \mu \mu')^{-1/2}, \quad \hat{B}_{\alpha} \equiv \int_0^{2\pi} d\beta' \delta(\beta' - \bar{\beta}) d\beta'$$

Here Ψ denotes the distribution function of these neutron collisions, α - an index which denotes the individual elements with the nuclear mass M_{α} occurring in the slowing down material, γ and β - the spherical angles of the vectors $\omega = \vec{p}/p$, \vec{p} - the momentum of the neutron, \vec{r} - radius vector of the neutron, $u = \ln(2mE_0/p^2)$, E_0 the initial energy of the neutrons, m - the mass of the

$\chi_{\text{un.eksp.1 teor.fis, 31, fasc.5, 893-895 (1956) CARD 2 / 5}$

PA - 1774

neutron, λ_{α} - the partial length of the free length of path of the neutron in consideration of the elastic collisions with nuclei with the mass M_{α} , λ - the total length of the free length of path of the neutron in the medium, m - the mass of the neutron. It further applies that:

$$f_{\alpha}(u) = \left[(M_{\alpha} + m) e^{-u/2} - (M_{\alpha} - m) e^{u/2} \right] / 2m$$

$$f_{\alpha_0}(u) = \left[(M_{\alpha} + m)^2 / 4mM_{\alpha} \right] e^{-u} \quad \text{or} \quad f_{\alpha_0}(u) = 0 \quad \text{at} \quad u \leq q_{\alpha} \quad \text{or} \quad u > q_{\alpha}.$$

$$q_{\alpha} = 2 \ln \left[(M_{\alpha} + m) / (M_{\alpha} - m) \right]$$

$$\mu_0 = \omega \omega' = \mu \mu' + \sqrt{1 - \mu^2} \sqrt{1 - \mu'^2} \cos(\beta' - \beta)$$

In the onedimensional problem Ψ is independent of β and consequently $\hat{B}_{\alpha} \Psi = \Psi$
It is then easy to develop K_{α} into a series according to LEGENDRE polynomials:

$$K_{\alpha} = \pi \sum_{l=0}^{\infty} (2l+1) P_l(\gamma_{\alpha}) P_l(\mu) P_l(\mu').$$

Žurn.eksp.i teor.fiz, 31, fasc.5, 893-895 (1956) CARD 3 / 5 PA - 1774

$K_\alpha(\mu, \mu', u)$ denotes the angular distribution (on the angle \mathcal{P}) in the laboratory system of the neutrons scattered by the nucleus. It is shown that even in the case of the scattering of neutrons by hydrogen nuclei the sum of some (n_0+1) initial terms of the series (which is here denoted by K_α^0) plays the main part. The remainder of the series $\tilde{K}_\alpha = K_\alpha - K_\alpha^0$ is written down in form of a finite expression. If all functions of u and μ are considered to be points of the metric space L_2 , the norm of the function $V(u, \mu)$ is defined as follows:

$$\|V\| = \int_0^u du e^{-u/2} \int_{-1}^1 d\mu V^2(u, \mu), \text{ and if the denotion } \varepsilon = \|\tilde{K}_\alpha\|^{1/2} / \|K_\alpha^0\|^{1/2}$$

is used, the value $\varepsilon = 0,33 < 1$ is found for $M = m$ and $n_0 = 1$.

Now n_0 is chosen in such a manner that 1. Instead of the punctiform kinetic equation $\hat{L}\Psi = \sum_\alpha \hat{K}_\alpha \hat{B}_\alpha \Psi + S$ the equation $\hat{L}\Psi^0 = \sum_\alpha \hat{K}_\alpha^0 \hat{B}_\alpha \Psi^0 + S$ is now investigated, where S denotes the density of the neutron sources and $\hat{L}_r(u, \mu, \beta) = \lambda(u)\omega \nabla + 1$. The influence exercised by \hat{K}_α can be looked upon as a perturbation. For this purpose one puts $\Psi = \Psi^0 + \Psi^1 + \Psi^2 +$

✓
 Žurn.eksp.i teor.fiz, 31, fasc.5, 893-895 (1956) CARD 4 / 5 PA - 1774
 + ... + $\Psi^{(n)}$ + By inserting this series into the aforementioned collision
 integral an equation for the determination of the n-th correction is obtained.
 The solutions of these equations can be written down with the help of a GREEN'S
 function G as follows: $\Psi^{(n)}(\vec{r}, u, \mu, \beta) = \int S^{(n-1)}(\vec{r}, u_1, \mu_1, \beta_1) \cdot$
 $G(\vec{r}, \vec{r}_1, u, u_1, \mu, \mu_1, \beta, \beta_1) d\vec{r}_1 du_1 d\mu_1 d\beta_1$, where $S^{(-1)} = S$. The aforementioned
 series converges towards $\Psi = \Psi^{(0)}$ and a formula is given for the error.

The GREEN'S function G satisfies the equation $\hat{L}\Psi^0 = \sum_{\alpha} \hat{K}_{\alpha}^0 \hat{B}_{\alpha} \Psi^{(0)} + S$, if one
 puts $S = \delta(r-r_1)\delta(u-u_1)\delta(\beta-\beta_1)$. If this equation is multiplied from the left
 with the operator L^{-1} and if GREEN'S function of the operator L is denoted
 with g, GREEN'S function can be put into the following shape in consideration
 of the boundary condition:

$$G = \sum_{\alpha} \left(d\vec{r}' \hat{K}_{\alpha}^0 \hat{B}_{\alpha} g(\vec{r}, \vec{r}', \mu, \beta) G(\vec{r}', \vec{r}_1, u', u_1, \mu', \mu_1, \beta', \beta_1) + \right. \\
\left. + g(\vec{r}, \vec{r}_1, u, \mu, \beta) \delta(u - u_1) \delta(\mu - \mu_1) \delta(\beta - \beta_1) \right).$$

The first part on the right side of this formula actually contains G^0 , the

Žurn.eksp.1 teor.fiz, 31, fasc.5, 893-895 (1956)

CARD 5 / 5

FA - 1774

sum of the first $\sum_{n=0}^{n_0} (2n+1)$ (in the onedimensional case of the (n_0+1))

spherical harmonics of the function G. Therefore determination of GREEN'S function is reduced to the determination of G^0 by the method of spherical harmonics, i.e. to the solution of a system with a small number of integral equations. These integral equations can be solved by taking the dependence of λ and $\lambda'_{\alpha(n)}$ on u into account. The GREEN'S function G as well as the functions $\Psi^{(0)}$ and $\Psi(n)$ are then determined in finite form. This is of particular importance for the computation of neutron distribution in media containing hydrogen (water, mineral oil, etc.), because the convergence of the usually used development of the distribution function according to spherical functions in this case is bad.

INSTITUTION: Institute for Geophysical Prospecting Methods.
Ministry for the Mineral Oil Industry of the USSR

71 211-230
TEMKIN, A.Ya.

Method for the approximate solution of the neutron kinetic equation.
Prikl. geofiz. no.17:211-230 '57. (MIRA 11:2)
(Neutrons) (Prospecting--Geophysical methods)
(Nuclear geophysics)

ZAPOROZHETS, V.M., kandidat tekhnicheskikh nauk; TEMKIN, A.Ya.

Using charged particle accelerators in investigating oil wells. Priroda 46 no.1:79-81 Ja '57. (MLBA 10:2)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut geofizicheskikh metodov razvedki Ministerstva neftyanoy promyshlennosti SSSR, Moskva (for Temkin).
(Petroleum research)

1. E. M. K. H. H. A.

3(5,6)

PHASE I BOOK EXPLORATION 304/2899

Vsesoyuzny nauchno-issledovatel'skiy institut geofizicheskikh metodov razvedki

Prikladnaya geofizika; sbornik statey, vyp. 23 (Applied Geophysics; Collection of Articles, No. 23) Moscow, Gostoptekhnizdat, 1959. 284 p. 3,500 copies printed.

Ed.: M.K. Polshkov; Exec. Ed.: M.M. Ruz'mina; Tech. Ed.: A. S. Polovina.

PURPOSE: This book is intended for scientific, engineering, and technical personnel of industrial geophysical exploration services.

COVERAGE: This is a collection of 14 articles by various authors on aspects of geophysical exploration. The material treated in the articles may be divided into four categories: the physical properties of rocks in specific geological regions; methods and techniques used in industrial geophysical exploration; concepts in the theory of electrical exploration, and the economics involved in geophysical operations. Specifically, the authors discuss the geologic structures of the central parts of the Russian Platform, southeastern Turbmenia, the West Siberian Plains, the eastern part of the Siberian Platform, and the Minusinsk basins; electrical frequency sounding, neutron logging, gamma spectrometry techniques, and the standard equipment and installations of the geophysical service of the petroleum industry in the USSR. References accompany each article.

ABSTRACTS: A.A. Density Characteristics of the Geological Profile of the Eastern Part of the Siberian Platform 112

Gelaktionov, A.B. Density of Sedimentary Beds of Ustyurt 127

Tarkov, A.P. Nature of the Anomalous Gravitational Field of the Minusinsk Basins 136

Teakin, A.Ya. Methods of Solving Problems in Neutron Logging 141

Kantor, I.I. The Effect of the Diameter of a Borehole on Instrument Readings in Neutron-Neutron Logging 174

Medentsev, G.A., P.M. Prokof'yev, A.I. Kholin, and A.P. Tsiforitskiy. Use of Differential Gamma-Spectrometry in Petroleum Geology 193

Voskobornik, M.Ya. The Speed of Electrical Logging in Combined Measurements With an Arbitrary Division of Channels 202

Polzakov, M.A. An Equivalent Electrical Schematic for an Electrode 217

Abb, E.A., V.M. Zaporozhets, B.I. Plotnikov, and L.A. Khatishvili. Some Problems in the Design of a Borehole Neutron Generator 226

Koilyev, P.T. Basic Assets of the Geophysical Services in the Petroleum Industry of the USSR 234

AVAILABLE: Library of Congress

Card 4/4

RU/58
12-31-59

16

TEMKIN, A.Ya.

Some methods of solving problems in the theory of neutron logging.
Prikl. geofiz. no.23:141-173 '59. (MIRA 13:1)
(Oil well logging, Radiation)

5 (4)

AUTHORS: Polak, L. S., Tonkin, A. P.

SCV/20-125-1-45/65

TITLE: On the Theory of Radiation Chemistry (K teorii radiatsionnoy khimii)

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 125, Nr 3, pp 534-537 (USSR)

ABSTRACT: The present paper deals with two problems: The taking into account of the interaction of the tracks and the method of approximate calculation of the quantity of the free radicals and final products of the (γ - and β -) radiolysis. The taking into account of the track structure, of its volume distribution and interaction is important for the investigation of the radiation-chemical reactions with low yields (~ 10 molecules per 100 ev). The reactions of chain character are by far less important. In the investigation of radiation-chemical reactions of non-chain character, (to which belong, for instance, the radiation cracking of hydrocarbons, some kinds of radiation polymerization, nitration, and many other reactions) high densities of the ionizing radiation must be applied in order to obtain a great yield of the final product per unit of time. High radiation densities,

Card 1/3

In the Theory of Radiation Chemistry

ISSN 0013-788X/86/0005-0006\$07.50/0

naturally, cause the interaction of the tracks. It is necessary, therefore, to take into account the mutual influence of the tracks of various ionizing particles if the density of the ionizing radiation is high. In this case the theory of radiation chemistry can be formulated as follows: it is necessary to find (for the instant of time t) the variation of the distribution of the ions and free radicals in the irradiated medium) which is caused by the formation of a track of given initial distribution of the ions and radicals in the instant of time $t_0 = t - t'$. An equation for the time

dependence of the density of the particles of the free radicals or ions is deduced. This equation, however, describes bimolecular reactions and does not take into account the possibility of secondary reactions. The authors then investigate some special cases: If a radiation of high density acts on polymers, there is nearly no diffusion of the polymer radical. In the case of reactions with yield of atomic hydrogen at high radiation densities the corresponding system of equations must contain an equation which describes the diffusion and various types of reactions of hydrogen atoms. The above-

Card 2/5

On the Theory of Radiation Chemistry

SOV/20-125-3-53/63

mentioned system of equations is then specialized and adapted for the case of low radiation densities, and the authors discuss a method for the approximate solution of this simplified system of equations. The formulae deduced then permit the computation of the number of free radicals and final products of the radiolysis. The method discussed in the present paper may be applied also to calculations by means of an electronic computer of discrete action. There are 3 references, 1 of which is Soviet.

ASSOCIATION: Institut neftekhimicheskogo sinteza Akademii nauk SSSR (Institute for Petroleum-chemical Synthesis of the Academy of Sciences USSR)

PRESENTED: December 11, 1958, by A. V. Tonchiyev, Academician

SUBMITTED: December 10, 1958

Card 3/3

TEMKIN, A.Ya.

Mechanism of the radiolysis of solid oxalic acid. Zhur. fiz.
khim. 34 no. 11:2503-2505 N '60. (MIRA 14:1)

1. Akademiya nauk SSSR, Institut neftekhimicheskogo sinteza.
(Oxalic acid) (Radiation)

86397

S.4300 1142, 1273, 1297
246820 1043

S/020/60/135/002/028/036
B004/B056

AUTHOR: Temkin, A. Ya.

TITLE: Theory of Thermalization of Hot Hydrogen Atoms and Their
Effect Upon the Dimer Yield in the Radiolysis of Alkanes

PERIODICAL: Doklady Akademii nauk SSSR, 1960, Vol. 135, No. 2,
pp. 373-376

TEXT: The present paper deals with the problem in which under the action of ionizing radiation upon gaseous organic substances part of the energy is converted into the kinetic energy of the fragments formed. In this case, kinetic energies may occur which are higher than the average energy of the thermal motion at the given temperature. In the radiolysis of alkanes, hot fragments, above all hot hydrogen atoms, will be formed. The action of these atoms upon the course of reaction is studied by an approximation method, the method of energy groups, such as is applied in the theory of slowing down and thermalization of neutrons. In the present case, i.e., the transition of hot H-atoms into thermal H-atoms, two kinds of particles

Card 1/5

186397

Theory of Thermalization of Hot Hydrogen
Atoms and Their Effect Upon the Dimer Yield
in the Radiolysis of Alkanes

S/020/60/135/002/028/036
B004/B056

are concerned, which may go over into each other. For a track process the system of equation is written which describes the diffusion, recombination, formation of free radicals, and thermalization of hot H-atoms:

$$\partial n_i / \partial t = D_i \Delta n_i - \sum_{j=1}^m a_{ij} n_i n_j - a_{iH} n_i n_H + A_i n_H + S_i \quad (1a);$$

$$\partial n_h / \partial t = D_h \Delta n_h - \Lambda n_h + S_h \quad (1b); \quad \partial n_H / \partial t = D_H \Delta n_H - \sum_{i=1}^m a_{iH} n_i n_H$$

$- a_{HH} n_H^2 + (\Lambda + \Lambda_H) n_H + S_H \quad (1c).$ D_i is the diffusion coefficient of i-type radicals; a is the recombination coefficient of i- and j-radicals; S_i is the number of i-radicals formed by the ionizing particle per unit time and volume; A_i is the macroscopic cross section for the formation of an i-radical by the hot H-atom; Λ is the macroscopic cross section for the collisions of the hot atom with gas molecules, in which the hot atom becomes thermal; n_i is the density of i-radicals. The subscript h refers

Card 2/5

86397

Theory of Thermalization of Hot Hydrogen
Atoms and Their Effect Upon the Dimer Yield
in the Radiolysis of Alkanes

S/020/60/135/002/028/036
B004/B056

to hot, and the subscript H to thermal hydrogen atoms. If the terms $a_{iH} n_i n_H$ are neglected, only the system of equations (1a), (1b) need be studied. The following example is given: At the instant t_0 of the formation of the track at the point with the radius vector \vec{r}_0 , hot atoms are formed. Within the sphere of radius R and with the same center, there are free radicals of a single type. The author develops a system of equations (2) for the dimer yield, using definitions taken from Ref. 3, which are not mentioned in this paper. If the intensity of radiation becomes so great that the tracks overlap and the reaction takes place within the entire volume, the group equations may be simplified. The following denotations are chosen: n_i : steady density of the hot atoms of energy E_i ; N: number of molecules of the initial substance per unit volume; σ_i : elastic scattering cross section of the hot atom of the i-group from a molecule, the atom going over into the (i+1) group; σ_{ik} , σ_{ic} : summational cross sections for all nonelastic collisions of a hot atom of the i-group, which cause its transition to the k-group ($k > i$). The following relations are found:

Card 3/5

86397

Theory of Thermalization of Hot Hydrogen
Atoms and Their Effect Upon the Dimer Yield
in the Radiolysis of Alkanes

S/020/60/135/002/028/036
B004/B056

$$N(\sigma_1 + \sum_{k=1}^n \sigma_{ik} + \sigma_{ic})n_1 = N[\sigma_{i-1}n_{i-1} \sum_{k=1}^{i-1} \sigma_{ik}n_k] + S_i (1 \leq i \leq m-1) \quad (3a);$$

$$N\sigma_{mc}n_m + a_{HH}n_m^2 + \sum_1 a_{H\mu}n_m n_{R,\mu} = N(\sigma_{m-1}n_{m-1} + \sum_{k=1}^{m-1} \sigma_{mk}n_k) + S_m \quad (3b).$$

Herefrom, the equations for the elementary events $\rho^{(\alpha)}$ are derived, e.g., for the reaction $H + M$ in which the H-atom vanishes, and for the reactions in which the hot atom passes from the i-group to the k-group. The author suggests to use this method of calculation when investigating the kinetics of processes in which hot molecules, radicals, or ions appear. He thanks L. S. Polak and N. Ya. Chernyak for discussions. There are 3 Soviet references.

Card 4/5

86397

Theory of Thermalization of Hot Hydrogen
Atoms and Their Effect Upon the Dimer Yield
in the Radiolysis of Alkanes

S/020/60/135/002/028/036
B004/B056

ASSOCIATION: Institut neftekhimicheskogo sinteza Akademii nauk SSSR
(Institute of Petrochemical Synthesis of the Academy of
Sciences USSR)

PRESENTED: June 3, 1960 by A. V. Topchiyev, Academician

SUBMITTED: June 3, 1960

Card 5/5

S/081/62/000/004/005/087
B149/B101

AUTHOR: Temkin, A. Ya.

TITLE: Accounting of the action of hot atoms and molecules in radiochemical kinetics

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 4, 1962, 73, abstract 4B508 (Tr. Tashkentsk. konferentsii po mirn. ispol'zovaniyu atomn. energii, v. I, 1959. Tashkent, AN UzSSR, 1961, 195-200)

TEXT: Equations were obtained describing the kinetics of chemical reactions induced by the passage of fast charged particles through a gas. The equations in general form take into account the diffusion and disintegration of the primarily excited molecules as well as the diffusion and recombination of the secondary radicals. In order to find the critical values of parameters at which irradiation induces a chain development of the process, the equations were linearized. Further simplification of the system associated with brief irradiation and preferential dissociation, rather than excitation of molecules is also considered. [Abstracter's note: Complete translation.]
Card 1/1

TEMKIN, A. YA.

16

PHASE I BOOK EXPLOITATION

SOV/6177

Akademiya nauk SSSR. Institut neftekhimicheskogo sinteza

Radioliz uglevodorodov; nekotoryye fiziko-khimicheskiye problemy
(Radiolysis of Hydrocarbons; Some Physicochemical Problems)
Moscow, Izd-vo AN SSSR, 1962. 207 p. Errata slip inserted.
5000 copies printed.

Resp. Eds.: A. V. Topchiyev, Academician, and L. S. Polak,
Doctor of Physics and Mathematics; Ed.: L. T. Bugayenko;
Tech Ed.: Ch. A. Zentsel'skaya.

PURPOSE: This book is intended for physical and industrial chemists
interested in the properties and behavior of irradiated hydro-
carbons.

COVERAGE: The book gives a systematic presentation of the results
of research on the radiolysis of hydrocarbons carried out from
1957 through 1961 at the Laboratory of Radiation Chemistry,
Institut neftekhimicheskogo sinteza AN SSSR (Institute of Petro-

Card 1/4

16

Radiolysis of Hydrocarbons (Cont.)

SOV/6177

chemical Synthesis, Academy of Sciences USSR). Although the results were obtained for individual compounds, they may be generalized and applied to other members of the same homologous series. The following persons participated in making the experiments and in writing the text: V. G. Beryozkin, V. E. Glushnev, Yu. A. Kolbanovskiy, I. M. Kustanovich, V. D. Popov, A. Ya. Tamkin, V. D. Timofeyev, N. Ya. Chernyak, V. A. Shakhray, E. B. Shlikhter, A. S. Shcherbakova, B. M. Negodov, A. Z. Peryshkina, M. M. Rytova, T. A. Tgina, Yu. B. Emin, A. M. Brodskiy, V. V. Voyevodskiy, P. Ya. Glazunov, B. A. Smirnova, and Yu. L. Khaik. References, mainly Soviet and English, follow individual chapters.

TABLE OF CONTENTS [Abridged]:

Foreword	3
Ch. I. Physicochemical Characteristics of Hydrocarbon Radiolysis	5
Card 2/4	

S/844/62/000/000/003/129
D290/D307

AUTHOR: Temkin, A. Ya.

TITLE: Application of the theory of multiple scattering of electrons in matter to the study of radiolysis mechanisms

SOURCE: Trudy II Vsesyuznogo soveshchaniya po radiatsionnoy khimii. Ed. by L. S. Polak. Moscow, Izd-vo AN SSSR, 1962, 34-37

TEXT: The case is considered of a source of fast electrons (energy E_0) that is uniformly distributed in an infinite homogeneous medium which only contains atoms of slow atomic number. The theory of multiple scattering of electrons in matter is used to show that the rates of production of singly-charged ions and excited molecules (S^+ and S^* respectively and the ratio S^*/S^+ all depend on E_0 . Considering the effect of the addition of a small quantity of inhibitor, it is shown that the inhibition coefficient also depends on E_0 . Che-

Card 1/2

Application of the theory ...

S/844/62/000/000/003/123
D290/D307

mical effects of radiation will depend on the energy spectrum of the radiation as well as on absorbed dose. The multiple scattering theory can be used in more general cases, e.g. in inhomogeneous media.

ASSOCIATION: Institut neftekhimicheskogo sinteza AN SSSR (Institute of Petrochemical Synthesis, AS USSR)

Card 2/2

43797

S/204/62/002/003/001/002
1032/1242

AUTHOR:

Temkin, A. Ya.

TITLE:

Contribution to the theory of the radiation polymerization of clathrate inclusion compounds

PERIODICAL: Neftekhimiya, v. 2, no. 3, 1962, 324-331

TEXT: The kinetics of radiation-induced polymerization of inclusion compounds is analysed theoretically. The dependence of the average molecular weight of the polymer on the duration of irradiation is calculated for two cases: (a) when the growth of the polymer chain proceeds from the point of initiation in both directions along the channel; and (b) when growth of the chain takes place in one direction only. Comparison with previously reported experimental results indicates that the growth of polymer chains from the point of initiation proceeds in one direction only, and that the observed decrease of average molecular weight of the polymer with duration of irradiation must be due to the fact that long channels, essential for growth of long polymer chains become more and more scarce as polymerization proceeds. In order to decide with greater certainty how polymerization proceeds, a small known amount of foreign molecules, that stop chain growth but are otherwise inactive, should be allowed during polymerization. Then the final average molecular weight of the

Card 1/2

Contribution to the theory of...

S/204/62/002/003/001/002
I032/I242

polymer will depend on the concentration of these inactive molecules in different ways, according to whether polymerization proceeds in both directions or in one only.

ASSOCIATION: Institut neftekhimicheskogo sinteza AN SSSR (Institute for Petrochemical Synthesis AS USSR) X

SUBMITTED: May 4, 1962

Card 2/2

GUBERMAN, Sh.A.; TEMKIN, A.Ya.

Two abstracts in "Referativnyi sbornik" (Series on petroleum). Izv.
AN SSSR. Ser. geofiz. no.6:951-952 Je 63. (MIRA 16:7)
(Oil well logging, Radiation)

BRODSKIY, A.M.; TEMKIN, A.Ya.

On the resonance theory of chemical reaction rates. Dokl. AN SSSR
152 no.1:127-130 S '63. (MIRA 16:9)

1. Institut neftekhimicheskogo sinteza AN SSSR. Predstavleno
akademikom Ya.B.Zel'dovichem.
(Chemical reaction, Rate of)

ALEKSANDROV, A.Yu.; DORFMAN, Ya.G.; LEPENDINA, O.L.; MITROFANOV, K.P.;
PLOTNIKOVA, M.V.; POLAK, L.S.; TEMKIN, A.Ya.; SHPINEL', V.S.

Resonance absorption spectra of γ -quanta and the magnetic
susceptibility of solutions of some organotin compounds.
Zhur. fiz. khim. 38 no.9:2190-2197 S '64. (MIRA 17:12)

1. Institut neftekhimicheskogo sinteza AN SSSR i Institut yadernoy
fiziki Moskovskogo gosudarstvennogo universiteta.

ABSTRACT: An attempt was made to observe the
reactions of hot atoms (atoms having very high kinetic energy, translation $\approx 10^6$ cm/
sec) with hydrocarbon molecules with the object of elucidating the mechanism of ionization.
Such collisions are common in radiation

"APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001755220006-3

APPROVED FOR RELEASE: 07/16/2001

CIA-RDP86-00513R001755220006-3"

TEMKIN, B.I.

"Study load and changes in the nervous activity of pupils during the school day."

report submitted at the 13th All-Union Congress of Hygienists, Epidemiologists and Infectionists, 1959.

TEMKIN, B. I.

Experience from teaching school hygiene in the pedagogic institute.
Gig. 1 san. no. 11:35-37 N '54. (MLRA 7:12)

1. Iz Kuybyshevskogo pedagogicheskogo instituta imeni V.V. Kuybysheva.
(HEALTH, education
in Russia, pedagogic institute)
(HYGIENE
school hyg. teaching in Russia, pedagogi, institute)

TEMKIN, B.I., dots.

Hygienic significance of workshop classes in daily school schedule
[with summary in English]. Gig. i san. 23 no. 8:34-36 Ag '58 (MIRA 11:9)

1. Iz Kuybyshevskogo pedagogicheskogo instituta imeni V.V. Kuybysheva.
(EDUCATION,
hyg. aspects of workshop classes (Rus))

TEMKIN, B.I.

Effect of work classes on the functional state of the nervous
system in school children. Uch. zap. Mosk. nauch.-issl. inst.
san. i gig. no.2:19-21 '59 (MIRA 16:11)

1. Kuybyshevskiy pedagogicheskiy institut imeni V.V.Kuybysheva.

*

TEMKIN, B.I.

Changes in the basal metabolism of schoolchildren during vacation
at Pioneer camps. *Pediatrics* 37 no.7:87 J1 '59. (MIRA 12:10)

1. Iz Kuybyshevskogo pedagogicheskogo instituta imeni Kuybysheva.
(METABOLISM)

TEMKIN, B.I., dotsent

Changes in the muscle tone in school children during work assignments. Gig.i san. 25 no.7:99-100 JI '60. (MIRA 14:5)

1. Iz Kuybyshevskogo pedagogicheskogo instituta imeni V.V. Kuybysheva.

(MUSCLES)

(SCHOOL CHILDREN)

TEMKIN, B.I., dotsent

Hygienic characteristics of the curriculum of schools with an
11-year course, and ways of making it more effective. *Pediatrics*
no.6:25-29 '61. (MIRA 14:9)

1. Iz Kuybyshevskogo pedagogicheskogo instituta imeni V.V.
Kuybysheva.
(SCHOOL HYGIENE)

TEMKIN, B.I., dotsent

Schedule and workload for pupils in the higher grades in
industrial education. Gig. 1 san. 26 no.7:40-45 J1 '61.
(MIRA 15:6)

1. Iz Kuybyshevskogo pedagogicheskogo instituta imeni
V.V. Kuybysheva.

(VOCATIONAL EDUCATION--HYGIENIC ASPECTS)

TEMKIN, B.I.

Changes in the higher nervous activity in students vacationing in
Pioneer camps. Vop.kur., fizioter.i lech.fiz.kul't. 27 no.2:128-
130 Mr-Apr '62. (MIRA 15:11)

1. Iz Kuybyshevskogo pedagogicheskogo instituta imeni V.V.
Kuybysheva.

(CONDITIONED RESPONSE) (CAMPING)

TEMKIN, B.I., dotsent

Functional body changes in boarding school pupils during lessons
at the school. *Pediatrica* 42 no.6:40-45 Je'63 (MIRA 17:1)

1. Kuybyshevskiy pedagogicheskiy institut imeni V.V.Kuybysheva.

A.E.S.

Geology

Dolomite flour of Shigulev origin. B. A. TANKIN.
Siber'skaya Press, 1940, No. 2, 9-10; *Chem. Zvest.*, 1940,
II, 1341; *Chem. Abs.*, 34, 4081 (1942).—A report on the
chemical composition and geologic conditions of a flourlike
dolomitic mass in a very extensive deposit of Shigulev is
presented. It contains R_2O , 0.09 to 1.8, CaO 30 to 32,
and Mg 20 to 21.8%. The material is so fine that it can
be used without further grinding in the glass industry.
It was tested and found a desirable raw material.

A.C. S.

G. H. H. H.

Reconstruction of spindle grinding-polishing tables SP-600. H. S. JAMES. *Soviet Army: Review from*, 1944, No. 7/8, pp. 13-16. — The movable parts of these polishing tables were quickly put out of commission because of the nature of their construction. This caused vibration and wobbling, which reflected adversely on the product. The gears, drives, bearings, etc., of the tables were completely redesigned, resulting in durable and uniform performance. M. Ho.

Use of slime-adulterated abrasives in polishing thick glass. H. S. TEMKIN. *Nekhod'sya i Keram. Pr.* 1944, No. 10-11, pp. 11-15. The following abrasives were used for polishing thick glass on Sh-1 machines: (1) corundum powders Nos. 140 and 325 and slime-adulterated 5-, 15-, 30-, and 60-min. powders, (2) corundum powders Nos. 200 and 325 and the slime-adulterated powders, and (3) corundum powders Nos. 200, 250, and 325 and the slime-adulterated powders. Best results were obtained with No. 3. No. 1 is preferred over No. 2 because it produced an even polish. The adulterated powders can be used without great harm to the process provided they are gradually and continually renewed so as not to allow much adulteration of the powders. Renovation of the adulterated powders with fresh powders should be made at least every 2 or 3 days. H.Z.K.

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

19

CA

Thermal stability of mirror reflectors. B. S. Tomkin. *Sokol'skiye i Karam. Prom.* 1945, No. 1-2, 7-10.—The existing method of testing the thermal stability of mirror reflectors by dipping in water at 65° followed by rapid immersion in water at 20° is not objective, does not indicate true thermal stability, and often results in the destruction of good products. The chem. compn. of glass of mirror reflectors, within the limits of ordinary cast mirror glass and Fourcault glass, has no noticeable effect on the thermal stability; hence, with regard to chem. compn., Fourcault glass can be considered satisfactory along with cast glass for making mirror reflectors. An objective method should be developed for testing the thermal stability of mirror reflectors which will approach the operating conditions at projector installations where the reflectors are protected from rapid cooling by means of a special glass. An app. should be devised to record the magnitude of residual stresses in the glass which could serve to det. the thermal stability without destroying the reflector.

B. Z. Kamich

COMMON SUBJECTS		PROCESSING AND PROPERTIES INDEX		19	
CA		<p>Protecting polished surfaces of glass prisms against alkali spotting. B. S. Tunkin and I. N. Burshstein. <i>Sokol'skiye i Keram. Prom.</i> 1943, No. 3, 8-10.—In polishing and grinding glass prisms, the use of alabaster to protect the polished surfaces resulted in the formation of spots requiring considerable polishing to remove. This defect was caused by the alkali in the alabaster and was facilitated by the high temp. of 66° attained during the process as well as by the length of the process (8-10 hrs.). This can be eliminated by acidifying the water used in prep., the alabaster and also by coating the polished surface with Iditol lacquer.</p> <p style="text-align: right;">B. Z. Kamich</p>			
MATERIALS INDEX		ASSOCIATED LITERATURE CLASSIFICATION		CITATION NUMBER	
FROM SYNOPTIC		TO TOPIC MAY GIVE CODE		FROM SOURCE	
SOURCE #A		SUBJECT OR KEY WORD		SUBJECT OR KEY WORD	
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100					

TENKIN, S. S. Cand. Tech. Sci.

Dissertation: "Peculiarities of the Effect of Chalk and Gypsum Slime on the Process of Glass Grinding." All-Union Sci Res Inst of Glass, 10 Mar 49.

CC: Vostochnaya Moskva, Mar, 1947 (Project #17006)

1st and 2nd process

PROCESSES AND PROPERTIES INDEX

5

C

BOOK

Grinding and Polishing of Glass (Shlifovka i polirovka stekla)
 B. S. TRAKIN. Published by Gosstroizdat, Moscow, 1947. 148
 pp. Price 3 60 rubles. Reviewed in *Steklo i Keram*, 5 (12) 22-23
 (1948). -In addition to the grinding and polishing of glass and
 the equipment used, much of the text is devoted to processes of
 glass manufacture. The book is replete with technical and typographical errors
 B. Z. K.

ASA-SLA METALLURGICAL LITERATURE CLASSIFICATION

1947-1948

1949-1950

1951-1952

1953-1954

1955-1956

1957-1958

1959-1960

1961-1962

1963-1964

1965-1966

1967-1968

1969-1970

1971-1972

1973-1974

1975-1976

1977-1978

1979-1980

1981-1982

1983-1984

1985-1986

1987-1988

1989-1990

1991-1992

1993-1994

1995-1996

1997-1998

1999-2000

2001-2002

2003-2004

2005-2006

2007-2008

2009-2010

2011-2012

2013-2014

2015-2016

2017-2018

2019-2020

2021-2022

2023-2024

2025-2026

2027-2028

2029-2030

2031-2032

2033-2034

2035-2036

2037-2038

2039-2040

2041-2042

2043-2044

2045-2046

2047-2048

2049-2050

2051-2052

2053-2054

2055-2056

2057-2058

2059-2060

2061-2062

2063-2064

2065-2066

2067-2068

2069-2070

2071-2072

2073-2074

2075-2076

2077-2078

2079-2080

2081-2082

2083-2084

2085-2086

2087-2088

2089-2090

2091-2092

2093-2094

2095-2096

2097-2098

2099-2100

2101-2102

2103-2104

2105-2106

2107-2108

2109-2110

2111-2112

2113-2114

2115-2116

2117-2118

2119-2120

2121-2122

2123-2124

2125-2126

2127-2128

2129-2130

2131-2132

2133-2134

2135-2136

2137-2138

2139-2140

2141-2142

2143-2144

2145-2146

2147-2148

2149-2150

2151-2152

2153-2154

2155-2156

2157-2158

2159-2160

2161-2162

2163-2164

2165-2166

2167-2168

2169-2170

2171-2172

2173-2174

2175-2176

2177-2178

2179-2180

2181-2182

2183-2184

2185-2186

2187-2188

2189-2190

2191-2192

2193-2194

2195-2196

2197-2198

2199-2200

2201-2202

2203-2204

2205-2206

2207-2208

2209-2210

2211-2212

2213-2214

2215-2216

2217-2218

2219-2220

2221-2222

2223-2224

2225-2226

2227-2228

2229-2230

2231-2232

2233-2234

2235-2236

2237-2238

2239-2240

2241-2242

2243-2244

2245-2246

2247-2248

2249-2250

2251-2252

2253-2254

2255-2256

2257-2258

2259-2260

2261-2262

2263-2264

2265-2266

2267-2268

2269-2270

2271-2272

2273-2274

2275-2276

2277-2278

2279-2280

2281-2282

2283-2284

2285-2286

2287-2288

2289-2290

2291-2292

2293-2294

2295-2296

2297-2298

2299-2300

2301-2302

2303-2304

2305-2306

2307-2308

2309-2310

2311-2312

2313-2314

2315-2316

2317-2318

2319-2320

2321-2322

2323-2324

2325-2326

2327-2328

2329-2330

2331-2332

2333-2334

2335-2336

2337-2338

2339-2340

2341-2342

2343-2344

2345-2346

2347-2348

2349-2350

2351-2352

2353-2354

2355-2356

2357-2358

2359-2360

2361-2362

2363-2364

2365-2366

2367-2368

2369-2370

2371-2372

2373-2374

2375-2376

2377-2378

2379-2380

2381-2382

2383-2384

2385-2386

2387-2388

2389-2390

2391-2392

2393-2394

2395-2396

2397-2398

2399-2400

2401-2402

2403-2404

2405-2406

2407-2408

2409-2410

2411-2412

2413-2414

2415-2416

2417-2418

2419-2420

2421-2422

2423-2424

2425-2426

2427-2428

2429-2430

2431-2432

2433-2434

2435-2436

2437-2438

2439-2440

2441-2442

2443-2444

2445-2446

2447-2448

2449-2450

2451-2452

2453-2454

2455-2456

2457-2458

2459-2460

2461-2462

2463-2464

2465-2466

2467-2468

2469-2470

2471-2472

2473-2474

2475-2476

2477-2478

2479-2480

2481-2482

2483-2484

2485-2486

2487-2488

2489-2490

2491-2492

2493-2494

2495-2496

2497-2498

2499-2500

2501-2502

2503-2504

2505-2506

2507-2508

2509-2510

2511-2512

2513-2514

2515-2516

2517-2518

2519-2520

2521-2522

2523-2524

2525-2526

2527-2528

2529-2530

2531-2532

2533-2534

2535-2536

2537-2538

2539-2540

2541-2542

2543-2544

2545-2546

2547-2548

2549-2550

2551-2552

2553-2554

2555-2556

2557-2558

2559-2560

2561-2562

2563-2564

2565-2566

2567-2568

2569-2570

2571-2572

2573-2574

2575-2576

2577-2578

2579-2580

2581-2582

2583-2584

2585-2586

2587-2588

2589-2590

2591-2592

2593-2594

2595-2596

2597-2598

2599-2600

2601-2602

2603-2604

2605-2606

2607-2608

2609-2610

2611-2612

2613-2614

2615-2616

2617-2618

2619-2620

2621-2622

2623-2624

2625-2626

2627-2628

2629-2630

2631-2632

2633-2634

2635-2636

2637-2638

2639-2640

2641-2642

2643-2644

2645-2646

2647-2648

2649-2650

2651-2652

2653-2654

2655-2656

2657-2658

2659-2660

2661-2662

2663-2664

2665-2666

2667-2668

2669-2670

2671-2672

2673-2674

2675-2676

2677-2678

2679-2680

2681-2682

2683-2684

2685-2686

2687-2688

2689-2690

2691-2692

2693-2694

2695-2696

2697-2698

2699-2700

2701-2702

2703-2704

2705-2706

2707-2708

2709-2710

2711-2712

2713-2714

2715-2716

2717-2718

2719-2720

2721-2722

2723-2724

2725-2726

2727-2728

2729-2730

2731-2732

2733-2734

2735-2736

2737-2738

2739-2740

2741-2742

2743-2744

2745-2746

2747-2748

2749-2750

2751-2752

2753-2754

2755-2756

2757-2758

2759-2760

2761-2762

2763-2764

2765-2766

2767-2768

2769-2770

2771-2772

2773-2774

2775-2776

2777-2778

2779-2780

2781-2782

2783-2784

2785-2786

2787-2788

2789-2790

2791-2792

2793-2794

2795-2796

2797-2798

2799-2800

2801-2802

2803-2804

2805-2806

2807-2808

2809-2810

2811-2812

2813-2814

2815-2816

2817-2818

2819-2820

2821-2822

2823-2824

2825-2826

2827-2828

2829-2830

2831-2832

2833-2834

2835-2836

2837-2838

2839-2840

2841-2842

2843-2844

2845-2846

2847-2848

2849-2850

2851-2852

2853-2854

2855-2856

2857-2858

2859-2860

2861-2862

2863-2864

2865-2866

2867-2868

2869-2870

2871-2872

2873-2874

2875-2876

2877-2878

2879-2880

2881-2882

2883-2884

2885-2886

2887-2888

2889-2890

2891-2892

2893-2894

2895-2896

2897-2898

2899-2900

2901-2902

2903-2904

2905-2906

2907-2908

2909-2910

2911-2912

2913-2914

2915-2916

2917-2918

2919-2920

2921-2922

2923-2924

2925-2926

2927-2928

2929-2930

2931-2932

2933-2934

2935-2936

2937-2938

2939-2940

2941-2942

2943-2944

2945-2946

2947-2948

2949-2950

2951-2952

2953-2954

2955-2956

2957-2958

2959-2960

2961-2962

2963-2964

2965-2966

2967-2968

2969-2970

2971-2972

2973-2974

2975-2976

2977-2978

2979-2980

2981-2982

2983-2984

2985-2986

2987-2988

2989-2990

2991-2992

2993-2994

2995-2996

2997-2998

2999-3000

C

PROCESSES AND PROPERTIES INDEX

Determination of the microgeometry of polished glass surfaces. B. S. TINKIN. *Nekhot'nyye i Keram. Prom.* 1947, No. 10, pp. 7-8. The following means are used in the Soviet Union for determining the microgeometry of polished glass surfaces: (1) Abbott profilometer, (2) Linnik interferometer, (3) Linnik pneumatic apparatus, and (4) optical contact method. The Linnik pneumatic apparatus is the most widely used; it gives an integral characteristic of the microgeometry of a small section of the polished surface but not of the depth of the separate "cavities." In the contact method, two pieces of glass are placed on an optical contact and polished (as one plate) along a surface perpendicular to the plane of the optical contact; then the optical contact is opened by heating, and the profile of the surface is measured under a microscope with the aid of an ocular micrometer. The Linnik interferometer is seldom used because it has a field of only 0.2 to 0.5 mm.² and, in addition, the procedure is complex. The Abbott profilometer is highly recommended.

H Z K

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

1947-1950

1951-1955

1956-1960

1961-1965

1966-1970

1971-1975

1976-1980

1981-1985

1986-1990

1991-1995

1996-2000

2001-2005

2006-2010

2011-2015

2016-2020

2021-2025

2026-2030

2031-2035

2036-2040

2041-2045

2046-2050

2051-2055

2056-2060

2061-2065

2066-2070

2071-2075

2076-2080

2081-2085

2086-2090

2091-2095

2096-2100

2101-2105

2106-2110

2111-2115

2116-2120

2121-2125

2126-2130

2131-2135

2136-2140

2141-2145

2146-2150

2151-2155

2156-2160

2161-2165

2166-2170

2171-2175

2176-2180

2181-2185

2186-2190

2191-2195

2196-2200

2201-2205

2206-2210

2211-2215

2216-2220

2221-2225

2226-2230

2231-2235

2236-2240

2241-2245

2246-2250

2251-2255

2256-2260

2261-2265

2266-2270

2271-2275

2276-2280

2281-2285

2286-2290

2291-2295

2296-2300

2301-2305

2306-2310

2311-2315

2316-2320

2321-2325

2326-2330

2331-2335

2336-2340

2341-2345

2346-2350

2351-2355

2356-2360

2361-2365

2366-2370

2371-2375

2376-2380

2381-2385

2386-2390

2391-2395

2396-2400

2401-2405

2406-2410

2411-2415

2416-2420

2421-2425

2426-2430

2431-2435

2436-2440

2441-2445

2446-2450

2451-2455

2456-2460

2461-2465

2466-2470

2471-2475

2476-2480

2481-2485

2486-2490

2491-2495

2496-2500

2501-2505

2506-2510

2511-2515

2516-2520

2521-2525

2526-2530

2531-2535

2536-2540

2541-2545

2546-2550

2551-2555

2556-2560

2561-2565

2566-2570

2571-2575

2576-2580

2581-2585

2586-2590

2591-2595

2596-2600

2601-2605

2606-2610

2611-2615

2616-2620

2621-2625

2626-2630

2631-2635

2636-2640

2641-2645

2646-2650

2651-2655

2656-2660

2661-2665

2666-2670

2671-2675

2676-2680

2681-2685

2686-2690

2691-2695

2696-2700

2701-2705

2706-2710

2711-2715

2716-2720

2721-2725

2726-2730

2731-2735

2736-2740

2741-2745

2746-2750

2751-2755

2756-2760

2761-2765

2766-2770

2771-2775

2776-2780

2781-2785

2786-2790

2791-2795

2796-2800

2801-2805

2806-2810

2811-2815

2816-2820

2821-2825

2826-2830

2831-2835

2836-2840

2841-2845

2846-2850

2851-2855

2856-2860

2861-2865

2866-2870

2871-2875

2876-2880

2881-2885

2886-2890

2891-2895

2896-2900

2901-2905

2906-2910

2911-2915

2916-2920

2921-2925

2926-2930

2931-2935

2936-2940

2941-2945

2946-2950

2951-2955

2956-2960

2961-2965

2966-2970

2971-2975

2976-2980

2981-2985

2986-2990

✓ Production of Polished Glass (*Proizvodstvo polirovanogo
stekla*) J. S. TERSTEN. Promstrolizdat, Moscow, 1959. 323
pp. Price R 14.00. Reviewed in *Seklo i Keram.*, 9 [10] 23-24
(1959). The book covers problems in making sheet glass, meth-
ods of casting and rolling, grinding and polishing of sheet glass,
laboratory control, polishing and grinding equipment, and
methods of calculation. The book reflects contemporary prog-
ress both in the Soviet Union and in foreign countries. B.Z.K.

POTOTSKAYA, G.V.; TEMKIN, B.S., nauchnyy redaktor; LITVAKOVSKIY, A.A.,
redaktor; DVORNIKOVA, N.I., tekhnicheskiy redaktor

[Production control in glass works manufacturing industrial glass]
Kontrol' produktsii na zavodakh tekhnicheskogo stekla. Moskva,
Gos. izd-vo lit-ry po stroit. materialam, 1953. 170 p.
[Microfilm] (MIRA 7:10)
(Glass manufacture)

TEMKIN, B.S.: kandidat technicheskikh nauk.

Ways for further development of the mirror industry. Leg.prom.
14 no.12:10-12 D '54. (MIRA 8:2)
(Mirrors)(Glass manufacture)

~~TRMKIN~~, B.S., kandidat tekhnicheskikh nauk; YUZHNIYA, Ye.A., redaktor;
MELNIKOVA, N.V., tekhnicheskii redaktor

[Mass production of mirrors] Proizvodstvo zerkal shirokogo potrebleniia.
Moskva, Gos. izd-vo mestnoi promyshlennosti RSFSR, 1956. 175 p.
(Mirrors) (MIRA 9:12)

TEMKIN, Boris Semenovich; GURENKOV, V.N., nauchnyy red.; GRINBERG, S.M.,
red.; GILSON, P.O., tekhn.red.

[Designing processes for assembly-line grinding and polishing
of sheet glass] Raschety protsessov konveiernogo shlifovaniia
i polirovaniia listovogo stekla. Moskva, Gos. izd-vo lit-ry po
stroit.materialam, 1957. 61 p. (MIRA 11:6)
(Grinding and polishing) (Glass)

TEMKIN, B.S.

Selecting methods for planning automatic flow lines in manufacturing polished glass. Stek. 1 ker. 17 no.6:19-22 Je '60.
(MIRA 13:6)

(Glass manufacture) (Conveying machinery)

KITAYGORODSKIY, I.I., doktor tekhn. nauk, prof.; KACHALOV, N.N., prof.;
VARGIN, V.V., doktor tekhn. nauk, prof.; YEVSTROP'YEV, K.S.,
doktor tekhn. nauk, prof.; GINZBURG, D.B., doktor tekhn. nauk,
prof.; ASLANOVA, M.S., doktor tekhn. nauk, prof.; GURFINKEL', I.Ye.,
inzh.; ZAK, A.P., kand. tekhn. nauk; KOTLYAR, A.Ye., inzh.; PAVLUSH-
KIN, N.M., doktor tekhn. nauk, prof.; Sentyurin, G.G., kand. tekhn.
nauk; SIL'VESTROVICH, S.I., kand. tekhn. nauk, dots.; SOLINOV, F.G.,
kand. tekhn. nauk; SOLOMIN, N.V., doktor tekhn. nauk, prof.; TEMKIN,
B.S., kand. tekhn. nauk; GLADYSHEVA, S.A., red. izd-va; TEMKINA, Ye.L.,
tekhn. red.

[Glass technology] Tekhnologiya stekla. Izd.3., perer. Moskva, Gos.
izd-vo lit-ry po stroit., arkhitekt. i stroit. materialam, 1961. 622 p.
(MIRA 14:10)

1. Chlen-korrespondent AN SSSR (for Kachalov).
(Glass manufacture)

TEMKIN, Boris Semenovich; KITAYGORODSKIY, I.I., doktor tekhn. nauk,
prof., retsenzent; NOVIKOVA, A.F., retsenzent; SULIMENKO, M.V.,
retsenzent; DUKHOVNIY, F.N., red.; SHAPENKOVA, T.A., tekhn.red.

[Technology of glass and glass products] Tekhnologiya stekla i
stekloizdelii. Moskva, Rostekhzdat, 1962. 458 p.
(MIRA 16:3)

(Glass)

BEREZHNOY, A.I.; BRODSKIY, Yu.A.; BRONSHTEYN, Z.I.; VEYNBERG, K.L.;
GALDINA, N.M.; GLETMAN, B.A.; GINZBURG, D.B.; GUTOP, V.G.;
GUREVICH, L.R.; DAUVAL'TER, A.N.; YEGOROVA, L.S.; KOTLYA,
A.Ye.; KUZYAK, V.A.; MAKAROV, A.V.; POLLYAK, V.V.; POPOVA,
E.M.; PRYANISHNIKOV, V.P.; Sentyurin, G.G.; SIL'VESTROVICH,
S.I., kand. tekhn. nauk, dots.; SOLOMIN, N.V.; TEMKIN, B.S.;
TYKACHINSKIY, I.D.; SHIGAYEVA, V.F.; SHLAIN, I.B.; EL'KIND,
G.A. [deceased]; KITAYGORODSKIY, I.I., zasl. deyatel' nauki i
tekhniki RSFSR, doktor tekhn. nauk, prof., red.; GOMOZOVA,
N.A., red.izd-va; KOMAROVSKAYA, L.A., tekhn. red.

[Handbook on glass manufacture] Spravochnik po proizvodstvu
stekla. [By] A.I.Berezhnoi i dr. Pod red. I.I.Kitaigorodskogo
i S.I.Sil'vestrovicha. Moskva, Gosstroizdat. Vol.2. 1963.
815 p. (MIRA 16:12)

(Glass manufacture)

TEMKIN, Boris Semenovich; BARKAN, Ye.Kh., retsenzents; IOFINOVA,
~~TS.B., red.~~

[Repair and manufacture of mirrors for consumers' use]
Remont i proizvodstvo bytovykh zerkal. Moskva, Legkaya
industriia, 1965. 117 p. (MIRA 18:4)

L 40318-66

ACC NR: AP6005335

arc gap, the control block contains a series-connected resistor-bridged capacitor, a resistor, and a unilaterally conducting element all connected in parallel to the arc gap, with the commutator input circuit connected in parallel to the element. A grid circuit and a thyatron cathode serve as the unilaterally conducting element (see Fig. 1). Orig. art. has: 1 figure.




Fig. 1. Device for activation and maintenance of an a-c welding arc.
1— capacitor; 2—resistor bridge; 3—resistor; 4— unilateral conducting element

[LD]

SUB CODE: 13/ SUBM DATE: 11Feb64

Card 2/2 MLP

BORISOV, V.T.; LYUBOV, B.Ya.; TIMKIN, D.Ye.

Calculation of the kinetics of solidification of metal ingots under various surface temperature conditions. Dokl.AN SSSR 104 no.2:223-226 S '55. (MLBA 9:2)

1. Institut metallovedeniya i fiziki metallov Tsentral'nogo nauchno-issledovatel'skogo instituta chernoy metallurgii. Predstavleno akademikom G.V. Kurdyumovym.
(Solidification) (Steel ingots)

TEMKIN, D.Ye.; TEUMIN, I.I.

Method of longitudinal oscillations for determining the
coefficient of internal viscosity. Zav. lab. 22 no.12: (MLRA 10:2)
1448-1451 '56.

1. Tsentral'nyy nauchno-issledovatel'skiy institut chernoy
metallurgii.
(Viscosity) (Deformations (Mechanics))

137-1958-2-2366

TEMKIN, D. YE.

Translation from: Referativnyy zhurnal. Metallurgiya, 1958. Nr 2. p 23 (USSR)

AUTHORS: Dobrovenskiy, V.V., Temkin D. Ye.

TITLE: How to Control Automatically the Growth of Single Crystals From a Melt by Means of Pulling With a Calculated Assignment of Input Parameters (K voprosu ob avtomaticheskoy regulirovani protsessy vyrashchivaniya monokristallov iz rasplava metodom vytyagivaniya s raschetnym zadaniyem vkhodnykh parametrov)

PERIODICAL: V sb.: Rost kristallov. Moscow, AN SSSR, 1957, pp 345-350

ABSTRACT: Solving the equation for the thermal conductivity of a crystal being grown from a melt following the Chochinskiy method, there being no radiation in the system of coordinates (which was stationary with respect to the solid phase), the A's arrived at the conclusion that the job of regulating an established crystallization process could be reduced to the task of controlling the temperature T_x of the upper end of a rod (the crystal) which is in contact with a cooler. A block diagram of an automatic control for T_x is given. The diagram indicates how the scheduled variations in the master e.m.f. were produced and how the difference between the e.m.f. of a thermocouple placed at the cooler and the master e.m.f. was transmitted

Card 1/2

137-1958-2-2366

How to Control Automatically the Growth of Single Crystals (cont.)

to a galvanometer; it shows also that the galvanometer was equipped with a photoelectric relay which was activated by any deviation, and it shows the amplifier and the actuating motor used to regulate the amount of water fed into the cooler.

Yu. Sh.

1. Crystals--Growing 2. Crystals--Thermal conductivity 3. Crystals--Mathematical analysis

Card 2/2